



Application	Dosage
Spreading into planting hole	<ul style="list-style-type: none">• 10 - 20 ml / plant (up to 12 cm pot)• up to 100 ml / plant (up to 10 l pot)• 20 ml / plant (up to 15 cm ø root ball)• up to 100 ml / plant (up to 40 cm ø root ball)
Existing plantings	<ul style="list-style-type: none">• up to 100 ml / plant in drilling holes• per 10 cm stem-ø 3 x 100 ml
Blending into substrat	<ul style="list-style-type: none">• 1 to 5 % (seedling production)• 5 to 10 % (plant production)
Blending with seeds	<ul style="list-style-type: none">• 10 – 20 ml / 1000 corn
Blending with fertilizer	<ul style="list-style-type: none">• 12 – 15 l / t
Blending with sowing at new grounds	<ul style="list-style-type: none">• up to 100 ml /m²
Surface application	<ul style="list-style-type: none">• up to 100 ml / m²
Transplantation bed	<ul style="list-style-type: none">• up to 100 ml / m furrow

For optimal development of the symbiosis please apply INOQ mycorrhiza products directly to the roots of the plants, either by spreading into the planting hole or distributing in the substrate/compost or soil surface.

These recommendations are not for INOQ Advantage concentrate.